

2024-03-27

Barriers and facilitators to parents' engagement with and perceived impact of a childhood obesity app: A mixed-methods study

Milne-Ives, M

<https://pearl.plymouth.ac.uk/handle/10026.1/22423>

10.1371/journal.pdig.0000481

PLOS Digital Health

Public Library of Science (PLoS)

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

RESEARCH ARTICLE

Barriers and facilitators to parents' engagement with and perceived impact of a childhood obesity app: A mixed-methods study

Madison Milne-Ives^{1,2}, Em Rahman³, Hannah Bradwell⁴, Rebecca Baines⁵, Timothy Boey⁶, Alison Potter³, Wendy Lawrence⁷, Michelle Helena van Velthoven⁸, Edward Meinert^{1,2,9,10*}



1 Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, United Kingdom, **2** Centre for Health Technology, School of Nursing and Midwifery, University of Plymouth, Plymouth United Kingdom, **3** South East School of Public Health, Workforce Training and Education Directorate, NHS England, United Kingdom, **4** School of Nursing and Midwifery, Faculty of Health, University of Plymouth, Plymouth, United Kingdom, **5** Peninsula Medical School, Faculty of Health, University of Plymouth, Plymouth, United Kingdom, **6** School of Medicine, University of Liverpool, Liverpool, United Kingdom, **7** Primary Care, Population Science and Medical Education, Faculty of Medicine, University of Southampton, Southampton, United Kingdom, **8** Nuffield Department of Primary Care Health Sciences, University of Oxford, Oxford, United Kingdom, **9** Department of Primary Care and Public Health, School of Public Health, Imperial College London, London, United Kingdom, **10** Faculty of Life Sciences and Medicine, King's College London, London, United Kingdom

* edward.meinert@newcastle.ac.uk

OPEN ACCESS

Citation: Milne-Ives M, Rahman E, Bradwell H, Baines R, Boey T, Potter A, et al. (2024) Barriers and facilitators to parents' engagement with and perceived impact of a childhood obesity app: A mixed-methods study. *PLOS Digit Health* 3(3): e0000481. <https://doi.org/10.1371/journal.pdig.0000481>

Editor: Haleh Ayatollahi, Iran University of Medical Sciences, IRAN (ISLAMIC REPUBLIC OF)

Received: November 9, 2023

Accepted: February 28, 2024

Published: March 27, 2024

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pdig.0000481>

Copyright: © 2024 Milne-Ives et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Deidentified individual participant data has been deposited in the Open Science Framework (OSF) repository

Abstract

Childhood obesity is a growing global health concern. Although mobile health apps have the potential to deliver behavioural interventions, their impact is commonly limited by a lack of sufficient engagement. The purpose of this study was to explore barriers and facilitators to engagement with a family-focused app and its perceived impact on motivation, self-efficacy, and behaviour. Parents with at least one child under 18 and healthcare professionals working with children were recruited; all participants were allocated to use the NoObesity app over a 6-month period. The mixed-methods design was based on the Non-adoption, Abandonment, Scale-Up, Spread, and Sustainability and Reach, Effectiveness, Adoption, Implementation, and Maintenance frameworks. Qualitative and quantitative data were gathered through semi-structured interviews, questionnaires, and app use data (logins and in-app self-reported data). 35 parents were included in the final analysis; quantitative results were analysed descriptively and thematic analysis was conducted on the qualitative data. Key barriers to engagement were boredom, forgetting, and usability issues and key barriers to potential impact on behaviours were accessibility, lack of motivation, and family characteristics. Novelty, gamification features, reminders, goal setting, progress monitoring and feedback, and suggestions for healthy foods and activities were key facilitators to engagement with the app and behaviours. A key observation was that intervention strategies could help address many motivation and capability barriers, but there was a gap in strategies addressing opportunity barriers. Without incorporating strategies that successfully mitigate barriers in all three determinants of behaviour, an intervention is unlikely to be successful. We

(DOI: [10.17605/OSF.IO/RUQ4G](https://doi.org/10.17605/OSF.IO/RUQ4G)). The dataset does not include the full demographic data collected to avoid indirect identification of participants. Following the guidance in Hrynaskiewicz et al. (2010), we have included only 3 indirect identifiers: age, gender, and ethnicity. Two participants did not consent for their data to be made available for future studies, so that data has been removed from the available dataset.

Funding: MMI and EM received research grant funding for this study from the former Health Education England, which is now the South East School of Public Health, Workforce Training and Education Directorate, NHS England (grant reference number: AM1000393). EM and MMI are supported by the NIHR Newcastle Biomedical Research Council (BRC). The views expressed in the paper belong to the authors and not necessarily those of the South East School of Public Health, NHS England, the University of Plymouth, the University of Liverpool, the University of Southampton, the University of Oxford, Newcastle University, the NIHR Newcastle BRC, or Imperial College London. The funding body had no editorial control and was not involved in the decision to submit the article for publication.

Competing interests: ER and AP are employees of the former Health Education England (now the South East School of Public Health, Workforce Training and Education Directorate, NHS England) and were involved in the development of the NoObesity apps. ER and AP contributed qualitative data and reviewed the final manuscript prior to submission, but the academic authors retained editorial control. MMI and EM received research grant funding for this study from the former Health Education England, which is now the South East School of Public Health, Workforce Training and Education Directorate, NHS England (grant reference number: AM1000393). EM and MMI are supported by the NIHR Newcastle Biomedical Research Council (BRC). The views expressed in the paper belong to the authors and not necessarily those of the South East School of Public Health, NHS England, the University of Plymouth, the University of Liverpool, the University of Southampton, the University of Oxford, Newcastle University, the NIHR Newcastle BRC, or Imperial College London. The funding body had no editorial control and was not involved in the decision to submit the article for publication. The other authors have declared that no competing interests exist.

highlight key recommendations for developers to consider when designing the features and implementation of digital health interventions.

Trial Registration: [ClinicalTrials.gov \(NCT05261555\)](https://clinicaltrials.gov/ct2/show/study/NCT05261555).

Author summary

Childhood obesity is a public health concern worldwide, but healthcare services lack the capacity to provide support and advice for all families on strategies for how it can be prevented or mitigated. This study explores what factors influence families' engagement with a mobile health app for childhood obesity prevention and their perceptions of its impact on their physical activity and eating behaviours. We found that novelty and interactivity—making the app fun and interesting—were key features that helped motivate families to keep using the app. Features of the app that enabled families set goals, keep track of their progress, and get suggestions for activities and healthy eating helped support their motivation and belief in their ability to engage in healthier behaviours. Beyond motivation, many families faced barriers related to their opportunity to engage in healthy behaviours such as lack of time, safe outdoor spaces, or affordable healthy food options. These findings highlighted the importance of understanding what prevents people from engaging with digital health interventions and the target behaviour change so that we can design interventions that help mitigate those barriers.

Introduction

Childhood obesity is a global concern affecting around 1 in 5 children worldwide [1]. Being overweight can negatively impact physical and mental health [2–8] and strain healthcare resources [9]. Determinants of childhood obesity are complex, including structural inequalities and genetics [10–12], but individuals can influence behavioural contributors like diet and activity [7,13–15]. The scope of the problem of childhood obesity in the UK—over a third of 10–11 year old children are overweight or obese [16]—requires support beyond the delivery capacity of clinical services. Mobile apps are a promising tool to support behaviour change because of their ubiquity [17–20], but their impact is commonly limited by low engagement [21–26].

Previous studies have supported the potential benefit of mobile apps for weight management for adults and children [18,27–33], especially for highly engaged users [27], but many interventions lack robust evidence of a long-term impact on weight [19,31,34–36]. It is essential that mobile health apps achieve sufficient user engagement to deliver impact [27,28,37]. Strategies for engaging users with digital interventions for childhood obesity are being examined [38], but further understanding of how to mitigate barriers to engagement is needed [39–42]. Behaviour Change Techniques (BCTs), the “smallest active ingredients” of interventions that can act to influence behaviour [43], are increasingly being incorporated into mobile health apps. Understanding how individual and contextual factors influence theoretical strategies, like BCTs, that aim to support engagement and behaviour change is necessary to optimise their behavioural and health impacts [44]. The purpose of this study was to gain insight into the context behind how and why parents engaged and disengaged with an app for childhood obesity, how it influenced their families' behaviour, and how potential improvements could be incorporated. Specifically, the study aimed to 1) generate barriers and facilitators that

influence engagement with the app and 2) to evaluate the app's impact on parents in terms of perceived motivation, self-efficacy, weight-related health behaviours, and communication with healthcare professionals (HCPs) [45].

Methods

Study design

We conducted a mixed-methods interventional study based on a Phase 1 implementation science design (Fig 1). Phase 1 studies focus on generating evidence around the implementation strategies and other factors that could influence acceptability, effectiveness, and successful adoption [46,47]. This is a critical step in the evaluative process, as it informs intervention refinement to improve impact in later efficacy- and effectiveness-oriented evaluations. The primary analysis examined qualitative data about barriers and facilitators to users' engagement with the app and its impact on motivation, self-efficacy, and behaviours. To improve the credibility of the qualitative results, they were triangulated with quantitative survey and app use data [45,48].

Three theoretical frameworks informed the study: the Reach Effectiveness Adoption Implementation Maintenance (RE-AIM) framework [49], the Non-adoption, Abandonment, Scale-up, Spread and Sustainability (NASSS) framework [50], and the Capability, Opportunity, Motivation—Behaviour (COM-B) model [51,52]. RE-AIM and NASSS informed the data collection plan, to capture a comprehensive set of individual- and system-level factors (Table 1 [49,50,53,54]). The COM-B model was used to analyse how various factors influenced engagement and behaviour [55–59]. The SRQR [60] (S1 Table) and TREND [61] checklists (S2 Table) were used to ensure completeness of reporting.

Intervention

The NoObesity app was developed by Health Education England (HEE) to prevent and manage childhood obesity. The term HEE is used as that was the organisation at the time of the study, but the organisation has since been incorporated into the South East School of Public Health, Workforce Training and Education Directorate, NHS England. The target audience included families wanting behavioural support (regardless of weight) and HCPs working with families [62]. It was not based on a particular behavioural theory but was co-produced with HCPs [62] and included features aligned with several BCTs (see S3 Table for a full list of




Fig 1. Study logic diagram.

<https://doi.org/10.1371/journal.pdig.0000481.g001>

Table 1. Alignment of RE-AIM and NASSS frameworks (based on [49,50,53,54]).

NASSS domains	RE-AIM dimensions	Key considerations	Key outcomes
Domain 1: The condition or illness Characteristics of users (clinical, socio-cultural, etc.)	N/A	<ul style="list-style-type: none"> Domain 1 emphasises the importance of understanding contextual factors and how they influence implementation 	<ul style="list-style-type: none"> Sample characteristics (demographics) Barriers and facilitators
Domain 2: The technology Features of technology associated with usability, acceptability, trustworthiness, and sustainability	Implementation Fidelity of intervention delivery (system), use of intervention strategies (user)	<ul style="list-style-type: none"> Both focus on key issues of usability, acceptability, and other factors associated with individual use RE-AIM captures issue of whether the intervention was delivered as intended 	<ul style="list-style-type: none"> System Usability Scale Key themes about usability, acceptability, and app features Barriers and facilitators Suggestions for improvement
Domain 3: The value proposition Evidence of impact and value, for users and suppliers (incl. safety, efficacy, benefit, affordability)	Effectiveness Impact on intended outcomes	<ul style="list-style-type: none"> Both emphasise the importance of evidence to support adoption of intervention (safety, positive impact on intended outcomes) NASSS considers impact of intervention for suppliers as well as users, and how that factors into the intervention's sustainability 	<ul style="list-style-type: none"> Key themes about motivation and perceived impact on self-efficacy and behaviours Ratings of app motivation strategies
	Adoption Number, proportion, and representativeness of places/people willing to adopt intervention (system)	<ul style="list-style-type: none"> RE-AIM 'adoption' also relates to the potential added value NASSS includes adoption by patients, staff, and setting, which is captured separately in RE-AIM ('Reach' and 'Adoption') Key questions around adoption: how much, how long, why, why not, what is required? 	<ul style="list-style-type: none"> Self-reported success at goals (in app and survey) Self-reported self-efficacy (in app and survey)
Domain 4: The adopter system Adoption (and continued use) by staff and users; reasons for non-adoption / abandonment; assumptions about use (capability, opportunity)	Reach Number, proportion, and representativeness participants (user)		<ul style="list-style-type: none"> Participation and drop-out data Key themes about adoption of HCP link and impact of name on adoption Ratings of HCP link and app name
Domain 5: The organization Organization's capacity and readiness for adoption and scale-up; recognition of need for and availability of dedicated resources; disruption to routines; work to implement	N/A	<ul style="list-style-type: none"> Domains 5 and 6 both related to various elements of the adoption and maintenance of the intervention, so have been analysed independently from the RE-AIM framework 	<ul style="list-style-type: none"> Key themes from qualitative interview with HEE representatives
Domain 6: The wider context Political, economic, regulatory, professional, sociocultural factors	N/A		
Domain 7: Embedding and adaptation over time Feasibility of ongoing adaptation; organisational resilience (willingness to reflect and adapt)	Maintenance Institutionalisation of intervention in routine processes (system), long-term impact on outcomes (user)	<ul style="list-style-type: none"> Both capture institutional / organisational maintenance RE-AIM also captures individual maintenance In this study, the organisational aspects of ongoing maintenance were closely aligned with key themes in Domains 5 and 6, and were grouped with them in the analysis 	<ul style="list-style-type: none"> Self-reported intention to continue use of app Self-reported length of use (if discontinued) App use data from system (length of use, most recent use date)

<https://doi.org/10.1371/journal.pdig.0000481.t001>

features, previously published in [45]). For families, the app helps them set behavioural goals to improve healthy eating and physical activity (e.g. cut down on snacks, walk to school, get the kids outside), track progress on their goals, and learn about healthy behaviours together through games and suggestions. For HCPs, the app provides training and tools for how to communicate effectively with families about childhood obesity (Fig 2). At the time of the study, the app was publicly available in the Apple App Store and Google Play store and promoted on HEE's website.

Sample and recruitment

From March to August 2020, a convenience sample of parents and HCPs living anywhere in the UK was recruited using social media advertising (Google Adwords, Instagram Ads) [63]. Participants self-selected to use the app in their daily lives over a 6-month intervention period




Fig 2. NoObesity Family and Professional app features.

<https://doi.org/10.1371/journal.pdig.0000481.g002>

based on the following inclusion criteria: (1) a parent/guardian (with at least one child any age under 18 years) OR a HCP working with families, (2) willing and able to use the app, (3) fluent in English, and (4) no previous use of the app. Individuals who were unable or unwilling to provide informed consent, had hearing impairments, or were known to the research team were excluded. Eligibility was not limited by weight to reflect real-world app use and the broad target audience. Informed consent was collected from all participants on enrolment using a Qualtrics survey. We also recruited two employees from HEE (ER, AP) who helped develop NoObesity to capture organisational-level factors influencing adoption and sustainability (Table 1).

Stratified random sampling (based on gender, ethnicity, and income) was used to invite a subset of 20 participants for semi-structured interviews (SSIs) from the 61 participants who completed the demographic questionnaire. Interviewed participants and those included in the quantitative data analysis (completed all questionnaires and passed an attention check question) received a £100 (USD\$139) Amazon voucher.

Procedure

Ethical approval was obtained from the University of Oxford Medical Sciences Interdivisional Research Ethics Committee (R62092/RE001) and the University of Plymouth's Faculty Research Ethics and Integrity Committee (19/20-1316). Thirty-minute SSIs were conducted on Microsoft Teams (S1 Text) and transcribed by a third-party company (Rev). To mitigate response bias, participants were informed that the interviewer was independent from HEE and had not been involved in intervention development. Three questionnaires were administered online using Qualtrics: consent, demographics based on OxIS 2019 [64] (September 2020), and the final questionnaire at the end of the 6-month intervention period (April 2021), which examined usability, use, and impact on motivation, self-efficacy, and behaviour and included attention check questions (S1 Text). App use data was collected in June 2021.

Outcomes

The primary outcome—barriers and facilitators to engagement and health behaviours—was explored through the qualitative SSIs. Secondary outcomes (Table 2) were examined through

Table 2. Quantitative secondary outcome measures.

Outcome	Outcome measure
Engagement with the app	App-captured dates of first and last logins
	Quantitative self-reported app use duration
Engagement with the study	Uptake (number of participants recruited)
	Dropout rates (number of participants completing each stage of the data collection)
Usability	System Usability Scale (SUS, scale: 0–100) [65]; completed by parents twice—once for themselves and once from their children's perspective
Motivation	Quantitative ratings of the impact of specific features on motivation (5-point Likert scale, see Table 3 for details)
Self-efficacy	Bandura's self-efficacy scale [66] (confidence in ability to do goal activities; scale: 0–100)
	In-app self-reported confidence in ability to achieve goals (6 star rating scale)
Perceived impact on behaviour	Quantitative questionnaire items about physical activity and healthy eating behaviours (5-point Likert scale, see Table 3 for details)
	In-app self-reported goal progress data
Opinions about app features	Quantitative questionnaire items about app features (5-point Likert scale, see Table 3 for details)

<https://doi.org/10.1371/journal.pdig.0000481.t002>

qualitative interviews, questionnaire, and app use data (including self-reported data and objective login data captured via the app admin portal).

Analysis

Qualitative data was analysed using a codebook thematic analysis approach [67,68] in Dedoose (version 9.0.17) [69]. To improve credibility—the alignment of participants' feedback with our representation of their feedback—four authors engaged with the data (first independently, then collaboratively) over a prolonged period. This ensured that individual preconceptions were challenged and alternative interpretations considered [70]. An initial list of codes was generated inductively by one author based on the transcripts, which were coded at the semantic level (interpreting only the explicit meaning of the participants' words) [71]. This codebook was provided to three independent authors who deductively coded the data (generating additional codes if needed). Codings then evolved through intensive collaboration between the authors. Themes and sub-themes were generated by one author as topic summaries [68] and then discussed and refined by all coders (S4 Table).

From the initial thematic framework, factors influencing engagement and perceived motivation, self-efficacy, and behaviours were deductively mapped to the COM-B model (S5 Table) [51,52,72]. A codebook for the mapping was created using the COM-B components [52] and the Theoretical Domains Framework (TDF) [73], which captures more detailed subthemes of the COM-B components (Table 3) and examples from the literature to guide coding in a digital health context (S6 Table) [58,59]. These factors are the main outcomes discussed. Quantitative survey and app use data were analysed using descriptive statistics. Where possible, these outcomes were triangulated with qualitative data.

Results

Sample characteristics

Consented participants included 225 parents/guardians and 6 HCPs. 61 parents completed the demographic survey; 15 of whom were interviewed. Almost 40% of parents (85/225) completed the final questionnaire, but only 16% (35/225) met the criteria to be included in the

Table 3. COM-B and TDF components and definitions [52,73].

COM-B	Definition [52]	TDF	Definition [73]
Physical capability	Physical skill, strength or stamina	Skills	An ability or proficiency acquired through practice
Psycho-logical capability	Knowledge or psychological skills, strength or stamina to engage in the necessary mental processes	Knowledge	An awareness of the existence of something
		Memory, attention, and decision processes	The ability to retain information, focus selectively on aspects of the environment and choose between two or more alternatives
		Behavioural regulation	Anything aimed at managing or changing objectively observed or measured actions
Physical opportunity	Opportunity afforded by the environment involving time, resources, locations, cues, physical 'affordance'	Environ-mental context and resources	Any circumstance of a person's situation or environment that discourages or encourages the development of skills and abilities, independence, social competence and adaptive behaviour
Social opportunity	Opportunity afforded by interpersonal influences, social cues and cultural norms that influence the way that we think about things, e.g. the words and concepts that make up our language	Social influences	Those interpersonal processes that can cause individuals to change their thoughts, feelings, or behaviours
Reflective motivation	Reflective processes involving plans (self-conscious intentions) and evaluations (beliefs about what is good and bad)	Social / professional role and identity	A coherent set of behaviours and displayed personal qualities of an individual in a social or work setting
		Beliefs about capabilities	Acceptance of the truth, reality or validity about an ability, talent or facility that a person can put to constructive use
		Optimism	The confidence that things will happen for the best or that desired goals will be attained
		Beliefs about consequences	Acceptance of the truth, reality, or validity about outcomes of a behaviour in a given situation
		Intentions	A conscious decision to perform a behaviour or a resolve to act in a certain way
		Goals	Mental representations of outcomes or end states that an individual wants to achieve
Automatic motivation	Automatic processes involving emotional reactions, desires (wants and needs), impulses, inhibitions, drive states and reflex responses	Reinforce-ment	Increasing the probability of a response by arranging a dependent relationship, or contingency, between the response and a given stimulus
		Emotion	A complex reaction pattern, involving experiential, behavioural, and physiological elements, by which the individual attempts to deal with a personally significant matter or event

<https://doi.org/10.1371/journal.pdig.0000481.t003>

quantitative analysis: completing demographic and final surveys and passing an attention check. No HCPs completed the study (Fig 3). Most participants were female and white, with a mean age of 39 years (SD 6.0; Table 4). Of the participants invited for interview via stratified random sampling, 8 responded and 5 completed the interviews. After a second round of stratified random sampling, all eligible participants were invited to reach a sample of 15 interviewed participants. Participants who did not complete follow-up assessments may still have been using the app, as it was publicly available and usage was not controlled.

Engagement with the intervention (adoption and use)

A large sample was initially recruited (n = 225), suggesting that there was interest in the app, but there was substantial drop-out from the study and discontinued app use (captured via self-report and app login data). Although most interviewed participants reported no plans to stop using NoObesity, only half of final survey respondents (19/35, 54%) reported still using the app and of the participants from whom app use data was collected (n = 26), only a third (9/26, 34%) had recently logged in by or after the end of the study (April 2021). On average, there were 5.1 months between account setup and most recent login (95% CI: 3.9–6.3, range: 0.0–




Fig 3. Participant flow diagram (HCPs: healthcare professionals; HEE: Health Education England).

<https://doi.org/10.1371/journal.pdig.0000481.g003>

10.4 months). This section examines the key capability, opportunity, and motivational factors identified from the thematic analysis as influencing engagement with the intervention (S5 Table). These could be grouped into two themes: motivation and usability (Figs 4–5).

Factors influencing motivation to engage

Most of the factors that influenced participants' motivation to engage related to the automatic and reflective motivation components of COM-B, although several overlapped with psychological capability (eg. TDF domain: 'memory, attention, decision processes, and knowledge') and physical opportunity (eg. TDF domain: 'environmental context' offered by the app). Novelty and variety were key for engagement and demonstrate this overlap; without something new to explore, participants felt bored, had reduced intentions to open the app, and needed increased cognitive effort to overcome these barriers (Box 1). Participants all valued feedback, but several found the app's feedback lacking or unhelpful. Visualisations of progress were suggested to help support motivation and more interactivity and gamification (e.g. daily

Box 1. Sample of participant quotes about factors affecting engagement with the app

"Once you've gone through the sections it has, then there's nothing else you can really do with it. So there's not much interaction . . . to keep you engaged." (ppt 2)

"Another reason why I didn't really do it is because. . . it feels quite repetitive having to click . . . Tuesday, Wednesday, Thursday, Friday [to record family progress] for every single challenge." (ppt 3)

"Had there been some new content on there, it would have encouraged me to come back more and more and go, "Oh, what more can I learn?" (ppt 7)

Table 4. Participant characteristics of the sample who completed the final quantitative survey (n = 35) and interviewed sample (n = 15).

Characteristic	Number (%) <i>final survey</i>	Number (%) <i>interviewed</i>
Gender		
Female	31 (89)	14 (93)
Male	4 (11)	1 (7)
Age (parent/guardian)		
30–39	20 (57)	11 (73)
40–49	12 (34)	3 (20)
50–59	3 (9)	1 (7)
Ethnicity		
White	32 (91)	12 (80)
Asian	2 (6)	1 (7)
Black	1 (3)	2 (13)*
Highest completed qualification		
Bachelor's degree or higher	23 (66)	8 (53)
Lower than Bachelor's degree	12 (34)	7 (47)
Annual household income before tax		
£12,500–19,999	3 (9)	1 (7)
£20,000–29,999	5 (14)	2 (13)
£30,000–39,999	7 (20)	5 (33)
£40,000–49,999	5 (14)	2 (13)
£50,000–59,999	4 (11)	2 (13)
£60,000–69,999	2 (6)	0 (0)
£70,000–79,999	1 (3)	0 (0)
≥ £80,000	6 (17)	1 (7)
Prefer not to say	2 (6)	2 (13)
Employment		
Working full time (30 hours a week or more)	12 (34)	3 (20)
Working part time (8–29 hours a week)	15 (43)	7 (47)
Doing housework, looking after children or other persons	4 (11)	3 (20)
Student	2 (6)	1 (7)
Permanently sick or disabled	2 (6)	1 (7)
Location		
A country village	5 (14)	2 (13)
A small city or town	24 (69)	10 (67)
The suburbs or outskirts of a big city	4 (11)	1 (7)
A big city	2 (6)	2 (13)
Number of adults in household		
One	3 (9)	2 (13)
Two	31 (89)	13 (87)
Three or more	1 (3)	0 (0)
Number of kids in household		
One	11 (31)	8 (53)
Two	16 (46)	5 (33)
Three	6 (17)	2 (13)
Four	2 (6)	0 (0)

*Not all interviewed participants completed the final quantitative survey

<https://doi.org/10.1371/journal.pdig.0000481.t004>




Fig 4. Summary of factors identified in the thematic analysis and how they were associated with engagement (Note: + and—signs refer to whether the factor had a positive or negative influence on the outcome).

<https://doi.org/10.1371/journal.pdig.0000481.g004>

challenges, tangible rewards, and friendly competition) to help engage children. Opinions about notifications were mixed: many participants appreciated the reminder, but several found them annoying or not delivered at appropriate times. A couple participants did not like the app's aesthetic and felt this hindered use.




Fig 5. Summary of key factors for engagement mapped to the COM-B components.

<https://doi.org/10.1371/journal.pdig.0000481.g005>

Positivity and a focus on behaviour rather than weight were important; some parents highlighted that while they might track calories or weight for themselves, they felt uncomfortable doing so for their children. This was also reflected in opinions of the name “NoObesity;” although most participants did not personally feel strongly about it, they thought it had negative connotations and that a more positive focus on health behaviours could improve new user engagement. The survey results supported this; participants reported somewhat disliking the name ($M = 2.6/5$, 95% CI: 2.2–3.0).

Factors associated with usability

Factors associated with ability to engage were primarily related to psychological capability and physical opportunity. Overall, interviewees found the app “quite easy to use,” in line with SUS scores ($M = 70.3/100$, 95% CI: 63.8–76.8), although ratings were slightly lower when parents completed the SUS from their child(ren)’s perspective ($M = 65.1/100$, 95% CI: 58.3–71.9). Barriers were primarily related to the technology (app freezing) and a lack of clarity and guidance in the app. Several parents reported not understanding the target audience, how to report goal progress as a family unit if only some of the family members had successfully achieved the goal, or how the app’s reward system worked, which hindered its potential impact on motivation (Box 2).

Box 2. Participant quotes about factors affecting usability

“Because of the iconography and how it looks, I think it’s quite easy to use. It’s quite self-explanatory.” (ppt 4)

“It could’ve possibly done with an introduction page to start with.” (ppt 6)

“There’s a lot of external links. I think you could quite easily get lost looking and looking for things.” (ppt 13)

“Two of us have gone out for a walk, but two of us haven’t. Do I score it [as having completed the family goal], do I not?” (ppt 11)

Perceived impact on behaviours

Perceived impact on health behaviours was mixed and often conflicting; most participants reported some progress on their goals but many said that the app did not have an impact on their behaviour (Box 3). In-app self-reported goal progress indicated that families were successful at achieving any particular goal for a mean of 7.5 weeks (95% CI: 5.5–9.5), a median of 3 weeks, and a mode of 0 weeks ($n = 27$). Factors relating to all three COM-B components were associated with the perceived impact on families’ motivation, self-efficacy, and health behaviours (Figs 6–7).

Facilitators for behaviours

Factors relating to perceived impact on motivation and self-efficacy were mostly related to motivation and psychological capability. Many participants found goal setting and progress recording motivating because it increased their awareness of their current behaviour and provided a sense of accountability (Box 4). Having “achievable, attainable” goals, “being able to

Box 3. Participant quotes about perceived impact of the app

“The healthier meals . . . last week we were able to tick off three or four, but I know we did more than that, it was just the actual ones I ticked off.” (ppt 15)

“[I] don’t necessarily think there were any changes. Let’s say, we live very close to school, so it’s easiest to walk. So that thing that I put in as our first goal about walking to school every day. We were going to use it every day, anyway.” (ppt 4)

input data,” and getting ideas for goal setting, physical activity, and healthy eating supported self-efficacy; which was largely aligned with the quantitative ratings of impact of app features on motivation, which were highest for ‘doing something together as a family’, ‘suggestions for activities and healthy eating’, and ‘goal setting’ (Table 5). Other facilitators included prompts and notifications and useful feedback (from app or a linked HCP). Median self-efficacy scores on the survey were around 70% (70/100) (range: 20–100%), similar to the mean confidence (out of 6 stars) users self-reported when setting goals on the app (3.9/6 stars, ~65%).




Fig 6. Summary of factors identified in the thematic analysis associated with the app’s perceived impact.

<https://doi.org/10.1371/journal.pdig.0000481.g006>




Fig 7. Summary of key factors for perceived impact mapped to the COM-B components.

<https://doi.org/10.1371/journal.pdig.0000481.g007>

Barriers to behaviours

Most behavioural barriers were associated with physical opportunity, including Covid-19 restrictions, time, weather, accessibility, and affordability (Box 5). There were also barriers relating to social opportunity (children’s willingness to try new foods and sibling teasing) and physical and psychological capability (barriers relating to children’s age and skills). Some participants suggested potential capability barriers, such as not knowing how to cook a healthy meal, but few experienced these barriers themselves. A lack of motivation was also mentioned; sometimes related to physical opportunity barriers (e.g. outdoor activity in bad weather), but not always with a reason given.

Box 4. Participant quotes about factors affecting motivation and self-efficacy

“This kind of brings it forward for you to actually acknowledge what you’re doing and what you’re not doing, and to help you to correct it. So it was like, I’m not going to have a snack today because I want to say that I managed to not have a snack today.” (ppt 2)

“I would have liked to be able to see trends. . . it’s nice to have a look at your graph, isn’t it?” (ppt 1)

“If you’ve downloaded it and it’s a family challenge, and it’s telling you to do it all together, it does make it easier, I think” (ppt 3)

“I thought the useful links were good . . . because if you’re not an outdoorsy family, you might not even know where to start” (ppt 8)

“Without [feedback from a HCP], it was completely lacking. [The app] didn’t give me feedback on whether my goals were good ideas. It didn’t give me feedback on how well I was doing meeting my goals.” (ppt 1)

Table 5. Participant ratings of the app and its features' perceived impact (n = 35).

Item	Mean (/5)	St. dev.	95% CI	Likert scale points meaning (1–5)
App's support for behaviour				
Set goals	3.5	1.4	3.0–4.0	Strongly disagree—strongly agree
Achieve goals	3.5	1.0	3.1–3.8	Strongly disagree—strongly agree
Eat healthier	3.4	1.3	3.0–3.8	Strongly disagree—strongly agree
Be more active	3.3	1.3	2.9–3.8	Strongly disagree—strongly agree
Perceived impact of app on motivation				
Eat healthier	2.7	1.1	2.3–3.0	Not at all—extremely effective
Be more active	2.6	1.1	2.2–3.0	Not at all—extremely effective
Perceived impact of motivation strategy				
Doing something together as a family	3.2	1.3	2.8–3.6	Not at all—extremely effective
Suggestions of activities	2.9	1.2	2.5–3.3	Not at all—extremely effective
Suggestions for healthier eating	2.8	1.0	2.5–3.2	Not at all—extremely effective
Goal-setting	2.8	1.3	2.4–3.3	Not at all—extremely effective
Self-monitoring	2.7	1.1	2.4–3.1	Not at all—extremely effective
Points / trophies ^a	2.7	1.3	2.2–3.1	Not at all—extremely effective
Notifications ^a	2.3	1.3	1.9–2.8	Not at all—extremely effective
Feedback about current behaviour	2.2	0.9	1.9–2.5	Not at all—extremely effective
Games	2.2	1.2	1.8–2.6	Not at all—extremely effective
Family photo	1.9	1.3	1.5–2.3	Not at all—extremely effective
Linking to a HCP	1.6	0.9	1.3–1.9	Not at all—extremely effective
Opinion of features				
Points / trophies	3.5	1.1	3.1–3.8	Dislike a great deal—like a great deal
Self-monitoring	2.7	1.0	2.4–3.1	Not at all—extremely useful
Information feedback	2.6	1.1	2.3–3.0	Not at all—extremely useful
Opinion on healthcare provider link				
Level of comfort	3.1	1.4	2.6–3.6	Extremely uncomfortable—extremely comfortable
Perceived usefulness	2.8	1.4	2.4–3.3	Not at all—extremely useful
Likelihood of linking	2.5	1.4	2.0–2.9	Extremely unlikely—extremely likely

^aSample was missing one data point (n = 34)

<https://doi.org/10.1371/journal.pdig.0000481.t005>

Box 5. Participant quotes about factors affecting behaviours

“We don't have outside space for us. So, it's an effort to do it every day.” (ppt 1)

“Some of the ideas were good, but there's just nowhere around here that we'd be able to do them.” (ppt 10)

“Our time is very limited. I go out to work and I'm not back till half five/six. So once we've had a family meal it's dark, the children aren't wanting to go back out. I'm not really wanting to go back out. It's cold.” (ppt 15)

“It's just easier to buy junk food, it's cheaper and easier to put. . . something in the oven rather than cook from scratch. And some people might not necessarily have the skills.” (ppt 8)

Perceived impact on communication

No HCPs completed the study and no participants connected with a HCP via the app (most had not seen a HCP during the intervention period because of Covid-19). Hypothetically, most parents felt comfortable communicating via the app and that the HCP link could provide useful feedback and accountability (Box 6), although a couple raised concerns about the app not reflecting true behaviour, bothering the HCP, discomfort discussing a sensitive topic, or data security.

Box 6. Participant quotes about factors affecting communication with HCPs

“I would have appreciated a real person saying, ‘well, that’s a good goal, or that’s a bit wishy-washy. Perhaps you could make that better’.” (ppt 1)

“It would be good because I think for me personally, if I’m accountable for something then, somebody checking in on you and going to look at what progress you’ve made. . . it would make me more determined to do it” (ppt 8)

“I don’t really feel I’ve got a relationship with my GP. So for me, I think I felt a bit reluctant about doing that just because I don’t know them.” (ppt 11)

Organisational factors

Interviews with HEE employees identified organisational-level factors that could influence the app’s sustainability. The app’s focus fits within UK priority areas around childhood obesity and digital-first healthcare. Its holistic approach, combining workforce development and service delivery, spans the mandate of several governmental bodies (since the interviews, HEE itself has been incorporated into NHS England) [74]. The HEE employees felt that this could be a benefit, as inter-agency collaboration could provide access to more expertise, but that it creates potential ambiguity around ownership, which could be a risk for longer-term maintenance.

Discussion

Main findings

This study used the COM-B model to explore factors associated with engagement and the perceived impact of a digital health app for childhood obesity and the NASSS and RE-AIM frameworks to generate insights for its implementation. Facilitators of engagement included novelty and variety, gamification and feedback, and a clear and positive tone. In terms of health behaviours, app features including BCTs [43] such as goal setting, problem solving (suggestions for goals), self-monitoring, feedback on behaviour, and instruction on how to perform the behaviour (suggestions for healthy foods and activities) helped support motivation and capability. Key barriers—including accessibility (e.g. Covid-19 restrictions, weather, affordability, and availability) and family characteristics—were largely related to opportunity, highlighting a gap in the app’s engagement and behaviour change strategies.

Recommendations in the context of existing evidence

There are many digital health interventions that have been developed and evaluated to target childhood obesity, but while studies often assess user perceptions and acceptability, there is

limited investigation of factors influencing engagement with mobile health apps for childhood obesity [31,75]. A recent evaluation of the Aim2Be app for childhood obesity identified different patterns of engagement and associated them with demographic characteristics, but not user perceptions [76], and another conducted a formative co-design process with parents to identify preferences for engagement and barriers to the health behaviours, which were largely similar to those identified here; however, these were not analysed in depth in that paper [77]. For this reason, we examine our findings in comparison to factors influencing engagement with digital health interventions more generally.

All three COM-B components are important for digital health intervention engagement and impact [78–85]. Facilitators for engagement—a simple visually-appealing interface, feedback and visualisations of progress, guidance, customisation, reminders, access to a HCP, and positive messaging—were aligned with previous recommendations in the literature [59,86–89]. Several factors, including common engagement strategies like gamification [90–92], notifications [82,93], and competition [94], could be barriers or facilitators. For example, in-app points and trophies alone may not be sufficiently meaningful rewards to support motivation [92] and notifications' impact can depend on their timing, frequency, and personalisation [82,93,95,96]. Friendly competition could help engage children by increasing fun and motivation [59]; however, parents were concerned about teasing and negative self-image. Social comparison can be demotivating [94], but this could be mitigated through team-based competition that de-emphasises the individual [42,97]. For childhood obesity, enabling intergroup family-team-based competition could help support engagement and focus on positive behaviours rather than weight.

Key facilitators for behavioural impact included awareness of behaviour, accountability, motivation, prompts, suggestions, family characteristics, and accessibility. Features that support a family focus [97], goal-setting, progress recording, positive messaging, feedback, and suggestions and inspiration could help support perceived motivation and self-efficacy in this context, in line with meta-analytic evidence in adults [98]. The relationship between motivation and accountability has been previously observed [99,100] and theoretically linked to adherence [101]. Accountability has the potential to help support behaviours not done for enjoyment [101,102], so more active involvement of a linked HCP with families via the app may have helped increase engagement with the intervention and behavioural goals; however, this may not work for everyone, as some participants worried that their app-reported progress did not reflect real behaviour and did not want to feel judged.

Key structural- and individual-level barriers were also largely aligned with existing evidence [81,85,89,103,104]. Structural inequities can hinder weight management and contribute to feelings of stigma or blame that reduce motivation [105,106]; mitigating these barriers (e.g. by highlighting affordable healthy meal options or locating free resources personalised to the users' location) will be necessary to enable positive health behaviours. Individual-level barriers were more aligned with motivation than capability [51], perhaps because families less aware of weight issues may have been less likely to use an app called “NoObesity.” As a key facilitator for engagement was the app's family focus, we recommend that developers consider the psychology behind their users' motivation to engage and how they can frame their intervention as something enjoyable rather than necessary but onerous [107]. Overall, we recommend that digital intervention designers research specific barriers (especially opportunity barriers) for their target populations and behaviours to identify the most appropriate means of mitigating them in that context. Table 6 highlights additional recommendations for intervention development, through the lens of the NASSS and RE-AIM implementation frameworks.

Table 6. Summary of implications for implementing similar interventions in the future.

NASSS domains	RE-AIM dimensions	Insights for designing interventions for successful implementation
Domain 1: The condition or illness	N/A	<ul style="list-style-type: none"> • Social determinants of health—such as income, environment, employment, etc.—are important factors that affect families' abilities to engage in healthy weight-related behaviours. • Digital health interventions should account for these contextual differences in users by incorporating advice and behavioural suggestions that meet their varying abilities and financial and environmental circumstances. • For example, this could include providing advice or recipes on how to eat healthy on a tight budget and including suggestions of physical activities that can be engaged in within a house if outdoor spaces are unavailable or inaccessible.
Domain 2: The technology	Implementation	<ul style="list-style-type: none"> • The importance of novelty as a factor to support engagement raises a potential issue for not-for-profit organisations that do not have capacity or funding for regular updates and uploading of new content for an intervention. • These types of limitations should be accounted for in the intervention design, for example, by setting up the intervention to provide content periodically over the course of the intervention or by enabling users to generate content (where appropriate). • Specific implementation strategies should be developed in collaboration with the target population, to ensure that the intervention meets their needs and circumstances.
Domain 3: The value proposition	Effectiveness	<ul style="list-style-type: none"> • To have a strong case to support public health interventions, evidence of positive impact is key—on an individual level, users should be able to see evidence of impact for themselves (eg. through progress monitoring and feedback from the intervention, or prompted self-reflection); on an organisational level, there needs to be significant evidence of positive impact to support widespread adoption. • Evaluations of effectiveness should include assessments of engagement as a prerequisite for impact and should include end-user perspectives on what key outcomes they consider desirable (eg. weight, mood, perceived energy, ability to do certain activities).
Domain 4: The adopter system	Adoption	<ul style="list-style-type: none"> • Given the high demand on HCPs' time and changes in practices and routines associated with implementation, perceived value, familiarity, and ease of use will be essential to clinical adoption [83]. • User-centred design processes should include a variety of clinical stakeholders, including those who would be directly engaging with the intervention and those responsible for managing adoption and implementation processes [108]. • This will be needed to identify and address challenges to adoption, whether they are related to perceived value and desirability of the intervention or structural barriers such as its lack of integration into existing routines
	Reach	<ul style="list-style-type: none"> • Social media and app store advertisement is unlikely to be sufficient to reach the target population in many cases; to increase reach, organisations should diversify dissemination by engaging with community groups and settings, specific to their target population's clinical and demographic characteristics. • The use of user-centred design with the target population could also improve reach by ensuring that, as well as being aware of the intervention, the target population expects that it will add value.
Domain 5: The organisation	N/A	<ul style="list-style-type: none"> • Within governmental and healthcare organisations, political changes could provide a potential challenge for continuity and responsibility of interventions; solutions for such issues will be case dependent, but should be considered from the outset to ensure there is a plan for sustainability and maintenance.
Domain 6: The wider context		
Domain 7: Embedding and adaptation over time	Maintenance	<ul style="list-style-type: none"> • Digital health interventions that aim to change behaviour contributing to health outcomes should use technical and social features that facilitate the incorporation of these behaviours into regular routines (see recommendations for features above).

<https://doi.org/10.1371/journal.pdig.0000481.t006>

Strengths, limitations, and future research

Strengths of this study included the use of mixed methods, theoretical frameworks (which guided the investigation of a comprehensive set of factors influencing adoption and implementation), and independent and collaborative coding by several authors for a rigorous thematic analysis. Due to time and resource limitations, mapping to the COM-B model was only completed by one author, which limits the robustness of that aspect of the analysis.

The main limitation was the sample; high dropout and non-representative demographics [109,110] created potential bias. Stratified random sampling was used to improve the diversity of the SSIs, but the limited overall sample and the need to invite all eligible participants to complete 15 interviews meant that interviewees' perspectives are likely to represent a particular set of lived experiences, reducing generalisability. The convenience method of sampling via social media also introduces potential bias, as participants who respond are likely to be those with

the strongest opinions about the intervention and the reach of the recruitment materials may not reflect a representative sample of the population. Random sampling could reduce this potential bias, although it may not reflect real-world engagement. Alternate methods of recruitment may have more success in increasing the sample size; for example, recruiting families and HCPs via schools, children's centres, primary care centres, and other community organisations. A larger-scale study with a more representative sample will be needed to evaluate the impact of the intervention on health behaviour change and childhood obesity. We also recommend that future studies capture demographic data immediately after consent, to enable analysis of participant characteristics in the initial and final samples.

Other limitations included the exclusion of children (to facilitate data collection and ethical approval) and the monetary reward for participants who completed the study (which might have incentivised app engagement). Data about the age of the children was not collected in the demographic data, but this was highlighted as an important factor in the qualitative analysis for consideration in future design and evaluation. Additional factors that could influence engagement and perceived impact are initial behavioural- and weight-related characteristics; these were not assessed as weight was not an inclusion criteria but may be an important factor to consider in future research.

The lack of data from HCPs meant we could not evaluate their perceptions of the intervention or their reasons for not participating in the study. The intervention period took place during the Covid-19 pandemic; unusually high demands on HCP's time might have precluded engagement with the study and intervention, but lack of participation could also be related to a lack of awareness of the study or HCP perceptions of the app. This will be an important area for future investigation, as previous research in digital health has suggested that integrating human support can improve engagement and is important for childhood obesity management [111,112].

Our ability to triangulate qualitative and quantitative data was also limited; the app use data that the system could record was relatively minimal and did not enable detailed examination of the frequency, intensity, time, or type of engagement (first and last login only, rather than more detailed data capturing number of logins or use of intervention features) [48]. The app is being redesigned based on theoretical frameworks and the findings of this study; this process will also improve its ability to capture app usage data. The limitations described here will be addressed in future larger-scale efficacy and effectiveness studies.

Conclusions

The growing prevalence of childhood obesity and the overwhelming demand for healthcare resources more generally has resulted in a need for easily accessible interventions to empower families to manage weight-related health. The benefits of digital health interventions are often limited by insufficient engagement, so understanding facilitators and barriers to engagement is essential. This paper highlights how the use of a theoretical behavioural framework can clarify key barriers to engagement with a digital health intervention and its target behaviour(s) and suggest mitigations. We recommend that digital intervention designers incorporate interactivity, novel content and suggestions, goal setting and progress monitoring, feedback and accountability, reminders, guidance on how to use the app, personalisation, and a positive and visually-appealing design. The caveat is that developers must identify whether these suggested intervention strategies align with engagement barriers and patient or other factors in their particular context; if strategies address capability and motivation barriers but not opportunity barriers, the intervention is unlikely to succeed. In terms of implementation, the NASSS and RE-AIM domains highlight key, interconnected factors that can influence the success of an

intervention—particularly important is the need to develop and demonstrate a value proposition that meets the needs and circumstances of the target users and clinical adopting systems, which can best be executed by adopting user-centred design practices to ensure that solutions to potential barriers are incorporated from early stages in development.

Supporting information

S1 Text. Semi-structured interview guide and study questionnaires.

(DOCX)

S1 Table. SRQR checklist.

(DOCX)

S2 Table. TREND checklist.

(DOC)

S3 Table. NoObesity Family and Professional apps functional overview (adapted from protocol) [45].

(DOCX)

S4 Table. Thematic framework.

(DOCX)

S5 Table. Thematic framework mapping factors to COM-B.

(DOCX)

S6 Table. COM-B and TDF coding framework with digital health examples.

(DOCX)

Author Contributions

Conceptualization: Em Rahman, Alison Potter, Wendy Lawrence, Michelle Helena van Velthoven, Edward Meinert.

Formal analysis: Madison Milne-Ives, Hannah Bradwell, Rebecca Baines, Timothy Boey.

Funding acquisition: Em Rahman.

Investigation: Madison Milne-Ives.

Methodology: Em Rahman, Alison Potter, Wendy Lawrence, Michelle Helena van Velthoven, Edward Meinert.

Supervision: Edward Meinert.

Visualization: Madison Milne-Ives.

Writing – original draft: Madison Milne-Ives.

Writing – review & editing: Madison Milne-Ives, Em Rahman, Hannah Bradwell, Rebecca Baines, Timothy Boey, Alison Potter, Wendy Lawrence, Michelle Helena van Velthoven, Edward Meinert.

References

1. Obesity and overweight. 9 Jun 2021 [cited 12 Apr 2022]. In: World Health Organisation [Internet]. Available: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
2. Morrison KM, Shin S, Tarnopolsky M, Taylor VH. Association of depression & health related quality of life with body composition in children and youth with obesity. *J Affect Disord.* 2015; 172: 18–23.

3. Beck AR. Psychosocial Aspects of Obesity. *NASN Sch Nurse*. 2016; 31: 23–27. <https://doi.org/10.1177/1942602X15619756> PMID: 26739931
4. Solmi F, Sharpe H, Gage SH, Maddock J, Lewis G, Patalay P. Changes in the Prevalence and Correlates of Weight-Control Behaviors and Weight Perception in Adolescents in the UK, 1986–2015. *JAMA Pediatr*. 2021; 175: 267–275. <https://doi.org/10.1001/jamapediatrics.2020.4746> PMID: 33196811
5. Pischon T, Boeing H, Hoffmann K, Bergmann M, Schulze MB, Overvad K, et al. General and abdominal adiposity and risk of death in Europe. *N Engl J Med*. 2008; 359: 2105–2120. <https://doi.org/10.1056/NEJMoa0801891> PMID: 19005195
6. Bass R, Eneli I. Severe childhood obesity: an under-recognised and growing health problem. *Postgrad Med J*. 2015; 91: 639–645. <https://doi.org/10.1136/postgradmedj-2014-133033> PMID: 26338983
7. Smith JD, Fu E, Kobayashi M. Prevention and Management of Childhood Obesity and its Psychological and Health Comorbidities. *Annu Rev Clin Psychol*. 2020; 16: 351. <https://doi.org/10.1146/annurev-clinpsy-100219-060201> PMID: 32097572
8. Djalalinia S, Qorbani M, Peykari N, Kelishadi R. Health impacts of Obesity. *Pak J Med Sci Q*. 2015; 31: 239–242. <https://doi.org/10.12669/pjms.311.7033> PMID: 25878654
9. Calculating the costs of the consequences of obesity. 2017 [cited 12 Apr 2022]. In: World Obesity Federation [Internet]. Available: <https://www.worldobesity.org/resources/resource-library/calculating-the-costs-of-the-consequences-of-obesity>
10. Robinson N, McKay JA, Pearce MS, Albani V, Wright CM, Adamson AJ, et al. The Biological and Social Determinants of Childhood Obesity: Comparison of 2 Cohorts 50 Years Apart. *J Pediatr*. 2021; 228: 138–146.e5.
11. Wu S, Ding Y, Wu F, Li R, Hu Y, Hou J, et al. Socio-economic position as an intervention against overweight and obesity in children: a systematic review and meta-analysis. *Sci Rep*. 2015; 5: 1–11. <https://doi.org/10.1038/srep11354> PMID: 26112253
12. Wang L, Morelen D, Alamian A. A prospective cohort study of the association between key family and individual factors and obesity status among youth. *Sci Rep*. 2022; 12: 1–10.
13. Preventing excess weight gain. NICE; 13 Mar 2015 [cited 28 Oct 2021]. In: National Institute for Health and Care Excellence [Internet]. Available: <https://www.nice.org.uk/guidance/ng7>
14. Strategies to Prevent & Manage Obesity. 9 Apr 2021 [cited 28 Oct 2021]. In: Centers for Disease Control and Prevention [Internet]. Available: <https://www.cdc.gov/obesity/strategies/index.html>
15. Ruth SM, Chan JW. Prevention of Overweight and Obesity: How Effective is the Current Public Health Approach. *Int J Environ Res Public Health*. 2010; 7: 765. <https://doi.org/10.3390/ijerph7030765> PMID: 20617002
16. Baker C. Obesity statistics. House of Commons Library; 2023 Jan. Available: <https://commonslibrary.parliament.uk/research-briefings/sn03336/#:~:text=Childhood%20obesity%20in%20England,and%20published%20by%20NHS%20Digital>.
17. Geirhos A, Stephan M, Wehrle M, Mack C, Messner E-M, Schmitt A, et al. Standardized evaluation of the quality and persuasiveness of mobile health applications for diabetes management. *Sci Rep*. 2022; 12: 1–10.
18. Wang Y, Xue H, Huang Y, Huang L, Zhang D. A Systematic Review of Application and Effectiveness of mHealth Interventions for Obesity and Diabetes Treatment and Self-Management. *Adv Nutr*. 2017; 8: 449–462. <https://doi.org/10.3945/an.116.014100> PMID: 28507010
19. Ghelani DP, Moran LJ, Johnson C, Mousa A, Naderpoor N. Mobile Apps for Weight Management: A Review of the Latest Evidence to Inform Practice. *Front Endocrinol*. 2020; 11. <https://doi.org/10.3389/fendo.2020.00412> PMID: 32670197
20. Rice L, Sara R. Updating the determinants of health model in the Information Age. *Health Promot Int*. 2019; 34: 1241–1249. <https://doi.org/10.1093/heapro/day064> PMID: 30212852
21. Meyerowitz-Katz G, Ravi S, Arnolda L, Feng X, Maberly G, Astell-Burt T. Rates of Attrition and Drop-out in App-Based Interventions for Chronic Disease: Systematic Review and Meta-Analysis. *J Med Internet Res*. 2020; 22: e20283. <https://doi.org/10.2196/20283> PMID: 32990635
22. Torous J, Lipschitz J, Ng M, Firth J. Dropout rates in clinical trials of smartphone apps for depressive symptoms: A systematic review and meta-analysis. *J Affect Disord*. 2020; 263: 413–419. <https://doi.org/10.1016/j.jad.2019.11.167> PMID: 31969272
23. Pratap A, Neto EC, Snyder P, Stepnowsky C, Elhadad N, Grant D, et al. Indicators of retention in remote digital health studies: a cross-study evaluation of 100,000 participants. *npj Digital Medicine*. 2020; 3: 1–10.
24. Birnbaum F, Lewis DM, Rosen R, Ranney ML. Patient engagement and the design of digital health. *Acad Emerg Med*. 2015; 22: 754. <https://doi.org/10.1111/acem.12692> PMID: 25997375

25. Yeager CM, Benight CC. If we build it, will they come? Issues of engagement with digital health interventions for trauma recovery. *Mhealth*. 2018; 4: 37. <https://doi.org/10.21037/mhealth.2018.08.04> PMID: 30363749
26. Baumel A, Muench F, Edan S, Kane JM. Objective User Engagement With Mental Health Apps: Systematic Search and Panel-Based Usage Analysis. *J Med Internet Res*. 2019; 21: e14567. <https://doi.org/10.2196/14567> PMID: 31573916
27. Dounavi K, Tsoumani O. Mobile Health Applications in Weight Management: A Systematic Literature Review. *Am J Prev Med*. 2019; 56. <https://doi.org/10.1016/j.amepre.2018.12.005> PMID: 31003801
28. Mclaughlin M, Delaney T, Hall A, Byaruhanga J, Mackie P, Grady A, et al. Associations Between Digital Health Intervention Engagement, Physical Activity, and Sedentary Behavior: Systematic Review and Meta-analysis. *J Med Internet Res*. 2021;23. <https://doi.org/10.2196/23180> PMID: 33605897
29. Oh C, Carducci B, Vaivada T, Bhutta ZA. Digital Interventions for Universal Health Promotion in Children and Adolescents: A Systematic Review. *Pediatrics*. 2022;149. <https://doi.org/10.1542/peds.2021-053852H> PMID: 35503335
30. Yau KW, Tang TS, Gorges M, Pinkney S, Kim AD, Kalia A, et al. Effectiveness of Mobile Apps in Promoting Healthy Behavior Changes and Preventing Obesity in Children: Systematic Review. *JMIR Pediatrics and Parenting*. 2022; 5. <https://doi.org/10.2196/34967> PMID: 35343908
31. Tully L, Burls A, Sorensen J, El-Moslemany R, O'Malley G. Mobile Health for Pediatric Weight Management: Systematic Scoping Review. *JMIR mHealth and uHealth*. 2020; 8: e16214. <https://doi.org/10.2196/16214> PMID: 32490849
32. Bonvicini L, Pingani I, Venturelli F, Patrignani N, Bassi MC, Broccoli S, et al. Effectiveness of mobile health interventions targeting parents to prevent and treat childhood Obesity: Systematic review. *Prev Med Rep*. 2022; 29: 101940. <https://doi.org/10.1016/j.pmedr.2022.101940> PMID: 36161123
33. Chai LK, Farletti R, Fathi L, Littlewood R. A Rapid Review of the Impact of Family-Based Digital Interventions for Obesity Prevention and Treatment on Obesity-Related Outcomes in Primary School-Aged Children. *Nutrients*. 2022; 14: 4837. <https://doi.org/10.3390/nu14224837> PMID: 36432522
34. Rowland SP, Fitzgerald JE, Holme T, Powell J, McGregor A. What is the clinical value of mHealth for patients? *npj Digital Medicine*. 2020; 3: 1–6.
35. Singh K, Drouin K, Newmark LP, Filkins M, Silvers E, Bain PA, et al. Patient-Facing Mobile Apps to Treat High-Need, High-Cost Populations: A Scoping Review. *JMIR Mhealth Uhealth*. 2016; 4: e136. <https://doi.org/10.2196/mhealth.6445> PMID: 27993761
36. Milne-Ives M, Lam C, De Cock C, Van Velthoven MH, Meinert E. Mobile Apps for Health Behavior Change in Physical Activity, Diet, Drug and Alcohol Use, and Mental Health: Systematic Review. *JMIR Mhealth Uhealth*. 2020; 8: e17046. <https://doi.org/10.2196/17046> PMID: 32186518
37. Yardley L, Spring BJ, Riper H, Morrison LG, Crane DH, Curtis K, et al. Understanding and Promoting Effective Engagement With Digital Behavior Change Interventions. *Am J Prev Med*. 2016; 51: 833–842. <https://doi.org/10.1016/j.amepre.2016.06.015> PMID: 27745683
38. Partridge SR, Redfern J. Strategies to Engage Adolescents in Digital Health Interventions for Obesity Prevention and Management. *Healthcare (Basel)*. 2018;6. <https://doi.org/10.3390/healthcare6030070> PMID: 29933550
39. Arthurs N, Tully L, O'Malley G, Browne S. Usability and Engagement Testing of mHealth Apps in Paediatric Obesity: A Narrative Review of Current Literature. *Int J Environ Res Public Health*. 2022;19. <https://doi.org/10.3390/ijerph19031453> PMID: 35162470
40. Henriksson P, Migueles JH, Söderström E, Sandborg J, Maddison R, Löf M. User engagement in relation to effectiveness of a digital lifestyle intervention (the HealthyMoms app) in pregnancy. *Sci Rep*. 2022; 12: 1–9.
41. Milne-Ives M, Homer S, Andrade J, Meinert E. Associations Between Behavior Change Techniques and Engagement With Mobile Health Apps: Protocol for a Systematic Review. *JMIR Res Protoc*. 2022; 11. <https://doi.org/10.2196/35172> PMID: 35348460
42. Milne-Ives M, Homer SR, Andrade J, Meinert E. Potential associations between behavior change techniques and engagement with mobile health apps: a systematic review. *Front Psychol*. 2023; 14: 1227443. <https://doi.org/10.3389/fpsyg.2023.1227443> PMID: 37794916
43. Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W, et al. The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Behav Med*. 2013; 46. <https://doi.org/10.1007/s12160-013-9486-6> PMID: 23512568
44. Greenwell K, Sereda M, Coulson NS, Geraghty AWA, Bradbury K, Hoare DJ. 'That's just how I am': a qualitative interview study to identify factors influencing engagement with a digital intervention for

- tinnitus self-management. *Br J Health Psychol.* 2021; 26: 727–747. <https://doi.org/10.1111/bjhp.12486> PMID: 33108049
45. Meinert E, Rahman E, Potter A, Lawrence W, Van Velthoven M. Acceptability and Usability of the Mobile Digital Health App NoObesity for Families and Health Care Professionals: Protocol for a Feasibility Study. *JMIR Res Protoc.* 2020; 9: e18068. <https://doi.org/10.2196/18068> PMID: 32706703
 46. Hull L, Goulding L, Khadjesari Z, Davis R, Healey A, Bakolis I, et al. Designing high-quality implementation research: development, application, feasibility and preliminary evaluation of the implementation science research development (ImpRes) tool and guide. *Implement Sci.* 2019; 14. <https://doi.org/10.1186/s13012-019-0897-z> PMID: 31412887
 47. Hamilton AB, Mittman BS. Implementation science in health care. In: Brownson, Colditz GA, Proctor EK, editors. *Dissemination and Implementation Research in Health: Translating Science to Practice* (2nd ed). Oxford University Press; 2017. <https://doi.org/10.1093/oso/9780190683214.003.0023>
 48. Short CE, DeSmet A, Woods C, Williams SL, Maher C, Middelweerd A, et al. Measuring Engagement in eHealth and mHealth Behavior Change Interventions: Viewpoint of Methodologies. *J Med Internet Res.* 2018;20. <https://doi.org/10.2196/jmir.9397> PMID: 30446482
 49. Glasgow RE, Vogt TM, Boles SM. Evaluating the public health impact of health promotion interventions: the RE-AIM framework. *Am J Public Health.* 1999; 89: 1322–1327. <https://doi.org/10.2105/ajph.89.9.1322> PMID: 10474547
 50. Greenhalgh T, Wherton J, Papoutsi C, Lynch J, Hughes G, A'Court C, et al. Beyond Adoption: A New Framework for Theorizing and Evaluating Nonadoption, Abandonment, and Challenges to the Scale-Up, Spread, and Sustainability of Health and Care Technologies. *J Med Internet Res.* 2017; 19: e367. <https://doi.org/10.2196/jmir.8775> PMID: 29092808
 51. Michie S, van Stralen MM, West R. The behaviour change wheel: A new method for characterising and designing behaviour change interventions. *Implement Sci.* 2011; 6: 42. <https://doi.org/10.1186/1748-5908-6-42> PMID: 21513547
 52. Michie S, Atkins L, West R. *The Behaviour Change Wheel: A Guide to Designing Interventions.* Silverback Publishing; 2014.
 53. What is RE-AIM? [cited 5 Nov 2021]. In: RE-AIM [Internet]. Available: <https://re-aim.org/learn/what-is-re-aim/>
 54. Glasgow RE, Harden SM, Gaglio B, Rabin B, Smith ML, Porter GC, et al. RE-AIM Planning and Evaluation Framework: Adapting to New Science and Practice With a 20-Year Review. *Front Public Health.* 2019; 7: 64. <https://doi.org/10.3389/fpubh.2019.00064> PMID: 30984733
 55. Huggins CE, Jong J, Leung GKW, Page S, Davis R, Bonham MP. Shift workers' perceptions and experiences of adhering to a nutrition intervention at night whilst working: a qualitative study. *Scientific Reports.* 2022; 1–10.
 56. Leather JZ, Keyworth C, Kapur N, Campbell SM, Armitage CJ. Examining drivers of self-harm guideline implementation by general practitioners: A qualitative analysis using the theoretical domains framework. *Br J Health Psychol.* 2022; 27: 1275–1295. <https://doi.org/10.1111/bjhp.12598> PMID: 35416355
 57. Keyworth C, Epton T, Goldthorpe J, Calam R, Armitage CJ. 'It's difficult, I think it's complicated': Health care professionals' barriers and enablers to providing opportunistic behaviour change interventions during routine medical consultations. *Br J Health Psychol.* 2019; 24: 571–592. <https://doi.org/10.1111/bjhp.12368> PMID: 30977291
 58. Szinay D, Jones A, Chadborn T, Brown J, Naughton F. Influences on the Uptake of and Engagement With Health and Well-Being Smartphone Apps: Systematic Review. *J Med Internet Res.* 2020; 22: e17572. <https://doi.org/10.2196/17572> PMID: 32348255
 59. Szinay D, Perski O, Jones A, Chadborn T, Brown J, Naughton F. Perceptions of Factors Influencing Engagement With Health and Well-being Apps in the United Kingdom: Qualitative Interview Study. *JMIR Mhealth Uhealth.* 2021; 9: e29098. <https://doi.org/10.2196/29098> PMID: 34927597
 60. O'Brien BC, Harris IB, Beckman TJ, Reed DA, Cook DA. Standards for reporting qualitative research: a synthesis of recommendations. *Acad Med.* 2014; 89: 1245–1251. <https://doi.org/10.1097/ACM.0000000000000388> PMID: 24979285
 61. Des Jarlais DC, Lyles C, Crepaz N. Improving the reporting quality of nonrandomized evaluations of behavioral and public health interventions: the TREND statement. *Am J Public Health.* 2004; 94. <https://doi.org/10.2105/ajph.94.3.361> PMID: 14998794
 62. King D, Rahman E, Potter A. NoObesity Apps—From Approach to Finished App. *Proceedings of the Future Technologies Conference (FTC) 2018.* 2019; 1145–1157.

63. NoObesity App Evaluation. 18 May 2020 [cited 25 Oct 2021]. In: NHS Health Education England South East—Working across Wessex [Internet]. Available: <https://wessex.hee.nhs.uk/wider-workforce/population-health/noobesity-digital-app/evaluation/>
64. Blank G. OxIS 2019 Questionnaire. All Parts. 2019 [cited 25 Oct 2021]. Available: <https://papers.ssrn.com/abstract=3522118>
65. Brooke J. SUS: A “Quick and Dirty” Usability Scale. *Usability Evaluation In Industry*. 1996; 207–212.
66. Bandura A. Guide for constructing self-efficacy scales. In: Urdan T, Pajares F, editors. *Self-Efficacy Beliefs of Adolescents*. Greenwich, CT: Information Age Publishing; 2006. pp. 307–337.
67. Brooks J, McCluskey S, Turley E, King N. The Utility of Template Analysis in Qualitative Psychology Research. *Qual Res Psychol*. 2015; 12: 202–222. <https://doi.org/10.1080/14780887.2014.955224> PMID: 27499705
68. Braun V, Clarke V. Toward good practice in thematic analysis: Avoiding common problems and be (com)ing a knowing researcher. *International Journal of Transgender Health*. 2022 [cited 24 May 2023]. <https://doi.org/10.1080/26895269.2022.2129597> PMID: 36713144
69. SocioCultural Research Consultants, LLC. Dedoose Version 9.0.17 web application for managing, analyzing, and presenting qualitative and mixed method research data. Los Angeles, CA; 2021. Available: www.dedoose.com
70. Nowell LS, Norris JM, White DE, Moules NJ. Thematic analysis: Striving to Meet the Trustworthiness Criteria. *Int J Qual Methods*. 2017; 16: 160940691773384.
71. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol*. 2006; 3: 77–101.
72. Attride-Stirling J. Thematic networks: an analytic tool for qualitative research. *Qual Res*. 2001; 1: 385–405.
73. Atkins L, Francis J, Islam R, O'Connor D, Patey A, Ivers N, et al. A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems. *Implement Sci*. 2017; 12: 1–18.
74. UKHSA operational launch—rebranding digital channels. 23 Sep 2021 [cited 4 Nov 2021]. In: UK Health Security Agency [Internet]. Available: <https://ukhsa.blog.gov.uk/2021/09/23/ukhsa-operational-launch-rebranding-digital-channels/>
75. Kouvari M, Karipidou M, Tsiampalis T, Mamalaki E, Poulimeneas D, Bathrellou E, et al. Digital Health Interventions for Weight Management in Children and Adolescents: Systematic Review and Meta-analysis. *J Med Internet Res*. 2022; 24: e30675. <https://doi.org/10.2196/30675> PMID: 35156934
76. González OD-J, Tugault-Lafleur CN, Jean Buckler E, Hamilton J, Ho J, Buchholz A, et al. The Aim2Be mHealth Intervention for Children With Overweight or Obesity and Their Parents: Person-Centered Analyses to Uncover Digital Phenotypes. *J Med Internet Res*. 2022;24. <https://doi.org/10.2196/35285> PMID: 35731547
77. Tripicchio GL, Kay M, Herring S, Cos T, Bresnahan C, Gartner D, et al. Development and Preliminary Feasibility of iByte4Health: A Mobile Health (mHealth) Pediatric Obesity Prevention Intervention to Engage Parents with Low-Income of Children 2–9 Years. *Nutrients*. 2021; 13. <https://doi.org/10.3390/nu13124240> PMID: 34959792
78. Borghouts J, Eikey E, Mark G, De Leon C, Schueller SM, Schneider M, et al. Barriers to and Facilitators of User Engagement With Digital Mental Health Interventions: Systematic Review. *J Med Internet Res*. 2021; 23: e24387.
79. König LM, Attig C, Franke T, Renner B. Barriers to and Facilitators for Using Nutrition Apps: Systematic Review and Conceptual Framework. *JMIR mHealth and uHealth*. 2021; 9: e20037. <https://doi.org/10.2196/20037> PMID: 34254938
80. Shabir H, D'Costa M, Mohiaddin Z, Moti Z, Rashid H, Sadowska D, et al. The Barriers and Facilitators to the Use of Lifestyle Apps: A Systematic Review of Qualitative Studies. *European Journal of Investigation in Health, Psychology and Education*. 2022; 12: 144–165. <https://doi.org/10.3390/ejihpe12020012> PMID: 35200235
81. Mauch CE, Laws RA, Prichard I, Maeder AJ, Wycherley TP, Golley RK. Commercially Available Apps to Support Healthy Family Meals: User Testing of App Utility, Acceptability, and Engagement. *JMIR Mhealth Uhealth*. 2021; 9: e22990. <https://doi.org/10.2196/22990> PMID: 33960951
82. Taki S, Russell CG, Lymer S, Laws R, Campbell K, Appleton J, et al. A Mixed Methods Study to Explore the Effects of Program Design Elements and Participant Characteristics on Parents' Engagement With an mHealth Program to Promote Healthy Infant Feeding: The Growing Healthy Program. *Front Endocrinol*. 2019; 10: 397. <https://doi.org/10.3389/fendo.2019.00397> PMID: 31293515
83. Thomas K, Neher M, Alexandrou C, Müssener U, Henriksson H, Löf M. Mobile phone-based lifestyle support for families with young children in primary health care (MINISTOP 2.0): Exploring behavioral

- change determinants for implementation using the COM-B model. *FrontHealth Serv.* 2022;0. <https://doi.org/10.3389/frhs.2022.951879> PMID: 36925820
84. Litterbach E-K, Russell CG, Taki S, Denney-Wilson E, Campbell KJ, Laws RA. Factors Influencing Engagement and Behavioral Determinants of Infant Feeding in an mHealth Program: Qualitative Evaluation of the Growing Healthy Program. *JMIR mHealth and uHealth.* 2017; 5: e8515.
 85. Willmott TJ, Pang B, Rundle-Thiele S. Capability, opportunity, and motivation: an across contexts empirical examination of the COM-B model. *BMC Public Health.* 2021; 21. <https://doi.org/10.1186/s12889-021-11019-w> PMID: 34051788
 86. Tang J, Abraham C, Stamp E, Greaves C. How can weight-loss app designers' best engage and support users? A qualitative investigation. *Br J Health Psychol.* 2015; 20: 151–171. <https://doi.org/10.1111/bjhp.12114> PMID: 25130682
 87. Hales S, Turner-McGrievy G, Fahim A, Freix A, Wilcox S, Davis RE, et al. A Mixed-Methods Approach to the Development, Refinement, and Pilot Testing of Social Networks for Improving Healthy Behaviors. *JMIR Human Factors.* 2016; 3: e8. <https://doi.org/10.2196/humanfactors.4512> PMID: 27026535
 88. Wei Y, Zheng P, Deng H, Wang X, Li X, Fu H. Design Features for Improving Mobile Health Intervention User Engagement: Systematic Review and Thematic Analysis. *J Med Internet Res.* 2020; 22: e21687. <https://doi.org/10.2196/21687> PMID: 33295292
 89. Taghizadeh S, Farhangi MA, Khodayari-Zarnaq R. Stakeholders perspectives of barriers and facilitators of childhood obesity prevention policies in Iran: A Delphi method study. *BMC Public Health.* 2021; 21: 2260. <https://doi.org/10.1186/s12889-021-12282-7> PMID: 34895191
 90. Tran S, Smith L, El-Den S, Carter S. The Use of Gamification and Incentives in Mobile Health Apps to Improve Medication Adherence: Scoping Review. *JMIR mHealth and uHealth.* 2022; 10. <https://doi.org/10.2196/30671> PMID: 35188475
 91. Lewis ZH, Swartz MC, Lyons EJ. What's the Point?: A Review of Reward Systems Implemented in Gamification Interventions. *Games Health J.* 2016; 5: 93–99. <https://doi.org/10.1089/g4h.2015.0078> PMID: 26812253
 92. Chow CY, Riantiningtyas RR, Kanstrup MB, Papavasileiou M, Liem GD, Olsen A. Can games change children's eating behaviour? A review of gamification and serious games. *Food Qual Prefer.* 2020; 80: 103823.
 93. Melcher J, Camacho E, Lagan S, Torous J. College student engagement with mental health apps: analysis of barriers to sustained use. *J Am Coll Health.* 2022; 70: 1819–1825. <https://doi.org/10.1080/07448481.2020.1825225> PMID: 33048626
 94. Tong HL, Laranjo L. The use of social features in mobile health interventions to promote physical activity: a systematic review. *npj Digital Medicine.* 2018; 1: 1–10.
 95. Bidargaddi N, Almirall D, Murphy S, Nahum-Shani I, Kovalcik M, Pituch T, et al. To Prompt or Not to Prompt? A Microrandomized Trial of Time-Varying Push Notifications to Increase Proximal Engagement With a Mobile Health App. *JMIR mHealth and uHealth.* 2018; 6: e10123. <https://doi.org/10.2196/10123> PMID: 30497999
 96. Freyne J, Yin J, Brindal E, Hendrie GA, Berkovsky S, Noakes M. Push Notifications in Diet Apps: Influencing Engagement Times and Tasks. *International Journal of Human–Computer Interaction.* 2017. <https://doi.org/10.1080/10447318.2017.1278896>
 97. Harris MA, Crone D. Motivations and barriers to engagement with a technology-enabled community wide physical activity intervention. *PLoS One.* 2020; 15: e0232317. <https://doi.org/10.1371/journal.pone.0232317> PMID: 32589658
 98. Samdal GB, Eide GE, Barth T, Williams G, Meland E. Effective behaviour change techniques for physical activity and healthy eating in overweight and obese adults; systematic review and meta-regression analyses. *Int J Behav Nutr Phys Act.* 2017; 14: 1–14.
 99. Bentley MR, Mitchell N, Sutton L, Backhouse SH. Sports nutritionists' perspectives on enablers and barriers to nutritional adherence in high performance sport: A qualitative analysis informed by the COM-B model and theoretical domains framework. *J Sports Sci.* 2019; 37: 2075–2085. <https://doi.org/10.1080/02640414.2019.1620989> PMID: 31124393
 100. Liddy C, Johnston S, Irving H, Nash K, Ward N. Improving awareness, accountability, and access through health coaching: qualitative study of patients' perspectives. *Can Fam Physician.* 2015; 61: e158–64. PMID: 25932483
 101. Oussedik E, Foy CG, Masicampo EJ, Kammrath LK, Anderson RE, Feldman SR. Accountability: a missing construct in models of adherence behavior and in clinical practice. *Patient Prefer Adherence.* 2017; 11: 1285. <https://doi.org/10.2147/PPA.S135895> PMID: 28794618

102. Deci EL, Koestner R, Ryan RM. A meta-analytic review of experiments examining the effects of extrinsic rewards on intrinsic motivation. *Psychol Bull.* 1999; 125: 627–68; discussion 692–700. <https://doi.org/10.1037/0033-2909.125.6.627> PMID: 10589297
103. Venkatesh A, Chang A, Green EA, Randall T, Gallagher R, Wildes JE, et al. Perceived Facilitators and Barriers to Engaging with a Digital Intervention among Those with Food Insecurity, Binge Eating, and Obesity. *Nutrients.* 2021; 13. <https://doi.org/10.3390/nu13072458> PMID: 34371967
104. Ray D, Sniehotta F, McColl E, Ells L. Barriers and facilitators to implementing practices for prevention of childhood obesity in primary care: A mixed methods systematic review. *Obes Rev.* 2022; 23: e13417. <https://doi.org/10.1111/obr.13417> PMID: 35064723
105. Brenton-Peters JM, Vallis M, Grant S, Consedine NS, Kirk SFL, Roy R, et al. Rethinking weight: Finding self-compassion for “weight management.” *Clin Obes.* 2023; 13: e12562.
106. Kebbe M, Perez A, Buchholz A, McHugh T-LF, Scott SS, Richard C, et al. Barriers and enablers for adopting lifestyle behavior changes in adolescents with obesity: A multi-centre, qualitative study. *PLoS One.* 2018; 13: e0209219. <https://doi.org/10.1371/journal.pone.0209219> PMID: 30562377
107. Van Kessel G, Kavanagh M, Maher C. A Qualitative Study to Examine Feasibility and Design of an Online Social Networking Intervention to Increase Physical Activity in Teenage Girls. *PLoS One.* 2016; 11: e0150817. <https://doi.org/10.1371/journal.pone.0150817> PMID: 26934191
108. Cornet VP, Daley C, Bolchini D, Toscos T, Mirro MJ, Holden RJ. Patient-centered Design Grounded in User and Clinical Realities: Towards Valid Digital Health. *Proceedings of the International Symposium on Human Factors and Ergonomics in Health Care.* 2019 [cited 6 Dec 2023]. <https://doi.org/10.1177/2327857919081023>
109. Analytical Impact Team. Overview of the UK population—Office for National Statistics. Office for National Statistics; 14 Jan 2021 [cited 5 Nov 2021]. Available: <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/articles/overviewoftheukpopulation/january2021>
110. Population of England and Wales. 1 Aug 2018 [cited 5 Nov 2021]. In: *Ethnicity Facts and Figures* [Internet]. Available: <https://www.ethnicity-facts-figures.service.gov.uk/uk-population-by-ethnicity/national-and-regional-populations/population-of-england-and-wales/latest>
111. Lipschitz JM, Pike CK, Hogan TP, Murphy SA, Burdick KE. The Engagement Problem: a Review of Engagement with Digital Mental Health Interventions and Recommendations for a Path Forward. *Current Treatment Options in Psychiatry.* 2023; 10: 119–135. <https://doi.org/10.1007/s40501-023-00297-3> PMID: 38390026
112. Wild CEK, Egli V, Rawiri NT, Willing EJ, Hofman PL, Anderson YC. “It’s more personal if you can have that contact with a person”: Qualitative study of health information preferences of parents and caregivers of children with obesity in New Zealand. *Health Soc Care Community.* 2022; 30. <https://doi.org/10.1111/hsc.13756> PMID: 35170827